



National Bee Unit

Hive Cleaning and Sterilisation

January 2022

Honey bee colonies are subject to infection or infestation by a range of pests and diseases. These include insects, mites, fungi, viruses, and bacteria, such as the microbes that cause American or European foulbrood (AFB and EFB) (*Paenibacillus larvae* and *Melissococcus plutonius*). Honey bees are social insects and are at risk of epidemics, so it is essential that beekeepers not only recognise the signs of such pests and diseases, but also know how to reduce their impact in colonies, apiaries and the locality. A key factor in preventing the spread of infection is good hygiene. The following Fact Sheet provides some advice about when and how you should be cleaning your hives and your equipment.

Be aware that to clean and sterilise your hives and equipment properly can be hard physical work. You need to make sure that you have all the necessary tools and equipment ready for the task before you get started. You also need to have appropriate protective clothing, including eye protection, strong waterproof gloves, steel capped boots etc. If possible, arrange for someone to help you on the day. A good place to get support and assistance can be through your local beekeeping association. Note that **not all treatments listed below are effective against foulbrood spores** and please bear in mind that all are potentially hazardous to human health. When using chemicals, always read the precautions on the manufacturers' label and take suitable safety precautions, especially when working with fire.

When should you clean and sterilise your beekeeping equipment?

It is good practice to clean and sterilise beekeeping equipment:

- When you bring it in from the field for storage (e.g. at the end of the season) or before reuse.
- When your colonies have experienced infection or infestation.
- Before using/reusing spare or empty hives, especially any second-hand items.
- When reusing any items which may have been associated with foulbrood infection.
- Whenever it is necessary to move items between colonies.

- Even apparently healthy colonies can harbour organisms that cause disease. The bacterium responsible for EFB and the spore forming bacterium that causes AFB can exist in a colony's combs for long periods of time, remaining capable of developing disease. It is therefore important to always maintain good standards of hygiene, even when pests and pathogens are not obviously present in an apiary. Replace old brood combs with foundation or new drawn comb on a regular basis and burn or render old comb.
- Always clean hive tools between inspections with a solution of washing soda. It is important to remember that it may not always be practical or safe to clean, disinfect and re-use beekeeping equipment once it has become contaminated, or is too old and/or damaged to withstand rigorous but necessary treatment.
- Methods for destruction are outlined in a separate section, below. Your local Bee Inspector will help and advise you in respect of hives and colonies that have been infected with notifiable diseases AFB and EFB.

Cleaning and disinfecting wooden hives

Dismantle the hive, and if possible, place all its component parts into a large domestic chest freezer (-20°C) for at least 48 hours before cleaning. This will kill insect pests, such as wax moth. Then place parts onto sheets of cardboard or newspaper inside an upturned roof to catch any debris that becomes dislodged as you work. Begin by scraping the boxes as thoroughly as possible, using a paint scraper, hive tool or other suitable instrument, to remove any adhering wax comb or propolis lumps.

Be especially careful when cleaning the internal corners of the boxes and the frame runners, as these provide ideal nooks and crannies that may harbour pests and pathogens. You may wish to consider removing "dirty" frame runners altogether and replacing them with new metal ones when you reassemble the disinfected hive. If you plan to sterilise your equipment by scorching (see below), remember to remove all plastic runners. During the process of scraping, bits of wax and propolis will fall onto your cardboard or newspaper sheets, and all this should subsequently be destroyed by burning when you have completed this part of the cleaning process. You will also need to clean your scraper before you put it away.

Once the wooden hive components are free of obvious dirt, they can be sterilised to kill germs. You can do this in several different ways:

Scorching with a blowlamp or handheld electric paint stripper

Fig. 1 below shows an example of scorching a brood box with a blow lamp to kill surface bugs. When doing this, use the tip of the 'blue flame' in the middle of the larger flame and make sure any remaining propolis boils. In addition, ensure that the timber darkens to a uniform coffee-brown colour, indicating that the wood has been heated to a sufficient temperature and for enough time to be properly sterilised. There is no need to burn the timber but be especially thorough in the corners.



Fig. 1: scorching a brood box with a blow lamp

Cleaning using washing soda crystals

Beekeeping tools and equipment (e.g. smokers and hive tools) can be cleaned using a solution of readily available and cheap washing soda crystals (Sodium Carbonate). This can be made up using 1 kg of washing soda to 4.5 litres of warm water with a dash of washing up liquid to help clean off propolis. Immerse the equipment in the solution, while using a wired brush, or similar tool to scrub off residues until the tools are clean.

Frame sterilisation can also be carried out in a similar manner. Lay some newspaper down on the surface where you will be working. Scrape all the frames as thoroughly as possible so that any excess propolis or wax falls onto newspaper underneath; this can then be burnt once you have finished. Prepare a mixture of soda solution (1:5) in a container such as an industrial water boiler (e.g. Burco boiler) large enough to hold honey frames and bring the water to the boil. Then submerge your frames in the boiling soda solution for a minute or until the frame is clean of any remaining wax and propolis. A scrub with a churn brush will help here. Remove the frames from the soda solution, rinse them down with water to remove residual Sodium Carbonate and then leave them to dry.

NB. This method is not suitable for sterilisation against American foulbrood.

Bulk-cleaning wooden hive parts in washing soda



Fig. 3: ensure that you are wearing appropriate protective clothing such as thick rubber gloves and safety goggles. Immerse the frames into the boiling soda solution for 1 to 2 minutes.



Fig. 4: once the equipment is clean, it can then be rinsed and left to dry, ready for reuse.

Additional equipment such as honey and brood boxes, queen excluders and crownboards can be cleaned in a soda solution in a similar way.

Chemical sterilisation with disinfectants

Brood boxes, supers and other beekeeping equipment can be effectively sterilised using disinfectants containing hypochlorite. Sodium hypochlorite is present at a concentration of about 3% in household bleach. Research has shown that immersion for twenty minutes in a solution of 0.5% sodium hypochlorite kills AFB spores and other bacteria. In this case you therefore need to make a solution of one part household bleach to five parts water. It is essential that the spores are in contact with the solution, so any items immersed must be thoroughly cleaned using the methods above first. For disinfecting one or two boxes, a suitable tray (such as one used for potting plants) could be used and each box wall treated individually, before rotating to treat the respective walls. For larger quantities of equipment, a deeper trough will be needed. After immersion in bleach, be sure to rinse the equipment down with water. Laboratory studies conducted at Fera Science Limited (Fera) suggest that other chemical disinfectants may be useful for general sterilisation, but further tests are required to ensure these are effective against foulbrood infection, especially AFB spores. When using this system **you must wear suitable protective clothing, protect your eyes and use rubber gloves.**

Chemical sterilisation with acetic acid

Combs can be sterilised to destroy the spores of chalkbrood, wax moth, and *Nosema spp.* disease of adult bees by using the evaporation fumes from acetic acid. **There is no evidence that this treatment is effective against AFB or EFB.** Acetic acid is available from chemical suppliers and online. Begin treatment by stacking the brood and/or super boxes containing combs to be sterilised on solid surface such as a board or solid hive floor. Note that acetic acid is corrosive and will attack metal and concrete. Vaseline on metal runners will help to protect them. It is also important to block off hive entrances, as acetic acid fumes are heavier than air and will travel from the top to the base of the stack, leaking out of any gaps or holes at the bottom. Place a non-metallic dish (saucer or similar container) on the top of the frames of the top box. Very carefully, put 80% acetic acid into the dish, allowing 120 ml acetic acid/box (e.g. 600 ml would therefore treat 5 boxes). Then, place an empty hive box on the top of the stack. Close the empty box on the top of the stack with a hive cover. Seal any joints between the boxes with wide adhesive tape to stop fumes escaping. Leave the stack for about one week to ensure sufficient fumigation. When the treatment is complete, the dishes of acid must be removed with caution and boxes should be thoroughly aired (at least two days) before they can be used again. When using this system **you must wear suitable protective clothing, protect your eyes and use rubber gloves.**

Boiling in caustic soda (Sodium hydroxide)

This treatment is not very practical for the average beekeeper due to the caustic nature of the bath and the problems of disposal of the used solution. It requires great care and caution. The method requires a bath of caustic soda solution, made up of 450 grams Sodium Hydroxide (available from chemical suppliers) in 38 litres of water, which must

then be brought to the boil. Wooden hive parts are then immersed in the solution for between 5 and 15 minutes, before being dipped in clean boiling water or washed with vinegar to neutralise the alkali. After drying, the parts will probably need re-nailing. When using this system, **you must wear suitable protective clothing, protect your eyes and use rubber gloves.**

Immersion into molten paraffin wax

Equipment dipped in paraffin wax must be heated to 160°C for 10 minutes. This is a dangerous process; and the flash point for paraffin wax is 199°C, so the temperature must be kept well below this level; **you will need specialist equipment and heavy-duty protective clothing**, which the average beekeeper does not have. For further details see a suitable beekeeping textbook or research online.

Irradiation

Wooden hive components such as stored supers and empty combs can be sterilised by gamma rays from a radioisotope of cobalt. This will kill infective organisms without damaging the equipment. Note: it is especially important that any equipment subjected to this treatment is entirely free from any signs of disease, such as the scales associated with AFB, prior to irradiation. This is because symptoms will remain indistinguishable visually from infectious untreated disease signs when the hive is put into use again and may lead to failure of diagnosis for a fresh infection. When using irradiation to sterilise hives after foulbrood infection, you should only do this under the guidance of your local Bee Inspector.

Recycling wooden frames




Fig. 5: steam cleaning apparatus

Hot Steam is very effective for melting wax from dirty wooden brood boxes and frames prior to their reuse. Apparatus consists of a source of steam, for example from a domestic wallpaper stripper, a plastic hose pipe, a collection tray, and a modified hive roof (Fig. 5). Equipment as described is available online.

Begin by placing a box filled with dirty frames onto the purpose-built collection tray, which needs to be similar in size and shape to a hive floor but with a small opening at one edge and cover with the lid. Attach the end of the hose pipe from the steam chamber into the top of the sealed box. As the wax softens and melts it runs down through the box and out of the opening in the tray, from where it can be collected for use in wax exchange (Fig. 6). The stripped and cleaned wooden frames can be reused. Allow at least one hour of treatment for each box of frames. When the steam cleaning process is complete, you can sterilise frames using washing soda treatment (see the 'Cleaning using washing soda crystals' section of this leaflet), before refitting with fresh foundation.



Fig. 6: collecting hot wax from a steam extractor

Cleaning and disinfecting plastic or polystyrene hive

Plastic beehives have been used in Europe for at least 30 years. Prior to sterilisation, plastic hives and components need to be carefully cleaned, just as you would a wooden hive – see information above. Once your plastic hive is clean, it is ready to be sterilised. However, options available are limited compared to the range of treatments you can use on wooden hives. Use of a blowtorch, for example, is clearly out of the question as this would simply melt the equipment, and the temperatures involved in paraffin wax treatment would also be inappropriate. Manufacturers of popular plastic hive brands recommend that the best available options are chemical disinfectants that contain hypochlorite or caustic soda, as described above for wooden hive sterilisation. Only hypochlorite-based products are recommended for AFB (see Dealing with EFB and AFB, below). Gamma irradiation is also effective. Any cleaning and treatment agent or solution should be tried on a small component or area to check that no damage will be caused.

Queen excluders

Queen excluders need to be cleaned and sterilised in different ways, depending on the type used. Begin by removing debris by scraping with a suitable tool (Fig. 7). It is easier to remove propolis when it is cold and wintery, as propolis will be brittle under these conditions. A wire brush is very useful to remove bits of wax and propolis. Wire excluders can then be scorched using a blowlamp, but if they are soldered be careful not to melt the solder joints as sometimes they are very 'soft'. If foulbrood has been present, zinc slotted excluders must be destroyed by burning. Otherwise, it is possible to scrub these clean with a solution of washing soda. This needs to be concentrated (1 kilogram of soda to 4.5 litres of water), and a dash of washing up liquid in the mix also helps. You will need to wear suitable protective clothing, protect your eyes and use rubber gloves. Plastic excluders can be disinfected in the same way as plastic hives and components, as described above.




Fig. 7: remove comb and debris, cleaning hive tool after use

How should I treat other equipment?

Hive tools can be cleaned in washing soda solution – ideally between each hive. Plastic equipment or components such as feeders and plastic frame ends can also be scrubbed off in a washing soda solution. Open-ended correx plastic sheets are difficult to clean inside so it is a good idea to seal up open ends prior to use. This does not apply to harbourage traps for monitoring for small hive beetle, which can be destroyed if necessary.

Cleaning beekeeping clothing

honey splashes and wiped off in a suitable manner. Beekeeping overalls should be washed regularly in the normal way. A small quantity of washing soda crystals mixed with the detergent helps to remove propolis. Make sure fencing style hoods are tucked inside and all zips are done up to avoid damage. Hats and veils with metal rings in cannot be washed in a machine.

Dealing with EFB and AFB

Both EFB and AFB are notifiable diseases. This means that if you suspect that your colonies are infected by either of these diseases you must report this to the National Bee Unit (NBU) who will provide confirmatory diagnosis. Details of how to do this are available on the NBU's BeeBase website (www.nationalbeeunit.com). Even before diagnosis is confirmed, close the hive and reduce the size of the entrance to one bee space, and take any other steps necessary to prevent the hive being robbed by other colonies in the area. Disinfect gloves and other beekeeping equipment with a strong solution of washing soda before examining other colonies. If AFB is subsequently confirmed, your local Bee Inspector will burn infected colonies, and hives must be sterilised before reuse.



Fig. 8: burning foulbrood-infected frames

Destruction of old, damaged and infected equipment

Wooden equipment can be burnt in a pit about 45 cm deep (Fig. 8), and the pit must be covered in afterwards to prevent any access to unburnt honey. Like any product, plastic hives become unserviceable because of damage or wear. When this occurs, they need to be disposed of safely. If the plastic has not been in contact with infectious diseases, then it may be suitable for recycling. Your Local Authority can give advice about this, which tends

to vary according to authority and region. Uninfected material can also be disposed of in landfill sites, but this facility may not be available in the future. Landfill is not an option for infectious material as it may remain exposed for some time. Additionally, when it becomes exposed in the future, it will become a source of infection for bees in the vicinity. Plastics must not be burnt except in a specialist plant. There are specialist companies that deal with the disposal of infected plastic materials, but they normally require plastics to be divided according to type before acceptance. They should be contacted for further information.

Are you registered on BeeBase?

BeeBase (www.nationalbeeunit.com) is the sophisticated IT system that supports the bee health programme. It includes custom designed databases comprising mapping, Geographical Information Systems (GIS) and spatial analysis capabilities. BeeBase contains all the apicultural information relating to the statutory bee health programme in England and Wales. It provides information on the activities of the NBU, legislation, honey bee pest and disease recognition and control, interactive maps, current research areas, and lots of helpful contacts. Many beekeepers find this an extremely useful resource and recognise the role that BeeBase has to play in the beekeeping community. If you register as a beekeeper on BeeBase you can request a free apiary visit from your local Bee Inspector, who can provide you with detailed help and advice on all aspects of beekeeping, including hive cleaning and sterilisation.

National Bee Unit

APHA, Room 11G03, York Biotech Campus,
Sand Hutton, York YO41 1LZ

Telephone: 03003030094 email: nbu@apha.gov.uk

Web site: www.nationalbeeunit.com

January 2022

© Crown copyright 2022

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ or email PSI@nationalarchives.gsi.gov.uk